

## Climate Projections and Impacts for the Republic of Palau

### Current Climate Changes

- Annual mean air temperatures have increased (~.6°C) in Palau since 1951.
- Interannual climate variability is high and is mainly influenced by the El Niño-Southern Oscillation (ENSO). Generally, El Niño years are drier than average and La Niña years are wetter.
- Sea levels have risen globally by 4-8 inches over the last 100 years; Tide gauge data in Palau indicates sea-level rise of ~4 inches since 1969 (University of Hawaii Sea Level Center 2015).
- Annual rainfall has increased ~3 inches (1948-2011) and is influenced by the West Pacific Monsoon (which brings heavy rainfall between June and August), the Intertropical Convergence Zone and Palau's location within the Pacific Warm Pool.
- Tropical cyclones are rare in Palau but can cause significant damage (e.g., Bopha in 2012, Haiyan in 2013). They affect Palau mainly between June and November.



### Climate changes over the next few generations

| Climate Feature        | Climate Projections  | Potential Impacts   |
|------------------------|--|---|
| <b>Sea Level Rise</b>  | Sea level is projected to increase in Palau by: <ul style="list-style-type: none"> <li>• ~5 inches over the next generation (by 2030)</li> <li>• ~10 inches by 2050</li> <li>• ~16-35 inches or more by the end of the century (2090 or four generations)</li> </ul> | Sea-level rise exacerbates flooding from high tides and storms. This can increase the potential for loss of lives, damage and loss of coastal homes, lands, and infrastructure, contaminated drinking water, and destruction of crops. Increased coastal erosion can result from higher sea levels especially when combined with large waves. Salinity intrusion can damage coastal aquifers and agricultural land. |
| <b>Air Temperature</b> | Annual temperatures will to continue to rise: <ul style="list-style-type: none"> <li>• ~0.8°C over the next generation (2030)</li> <li>• ~1-2°C by 2050</li> <li>• ~2-4°C by 2090</li> </ul>   | Impacts to human health and health systems related to heat stress if working outside or outdoor recreation. Increased need for cooling systems and energy required for cooling. Air temperature impacts sea surface temperature, storms, and precipitation, also impacting agriculture and water resources.   |
| <b>Storm Patterns</b>  | Globally, cyclones are projected to be fewer but the ones that occur will be of higher intensity. Models show inconsistent results for Palau; most suggest a decrease in formation (20-50%), but the confidence for these projections is low.                        | More severe cyclones when they do occur and combined with sea level rise will result in increased flooding and potentially coastal change resulting in damage and loss of lives, coastal homes, land, and infrastructure.   |

## Climate change impacts and natural hazards in Palau



Coastal erosion and typhoon impacts in Palau (Source: EU-SOPAC (EDF9) Project Report 152 2009; Office of the President, Republic of Palau)

| Climate Feature                      | Climate Projections   | Potential Impacts   |
|--------------------------------------|---|---|
| <b>Sea-Surface Temperature (SST)</b> | <p>Projected increases in SST:</p> <ul style="list-style-type: none"> <li>• ~0.8°C over the next generation (2030)</li> <li>• ~1-2°C by 2050</li> <li>• ~2-4°C by 2090</li> </ul>   | <p>Coral bleaching is expected to increase. When sea temperatures increase 1–2°C above the normal maximum for &gt; 4-6 week, coral bleaching is likely. Coral diseases may also increase due to warming seas. Coral bleaching and disease can adversely affect reef-dependent species and reduce services reefs provide (tourism; coastal protection; food/livelihoods; habitat; medicine).</p>   |
| <b>Rainfall Patterns</b>             | <p>Average rainfall is projected to increase, especially in the wet season, along with more extreme rain events (~2% increase by 2030; ~4% increase by 2050; ~8% increase by 2090). Droughts are projected to decline in frequency. There will still be wet and dry years and decades due to ENSO-related variability, but most models show that the long-term average is expected to be wetter.</p>  | <p>Increases in rainfall intensity will lead to increasing flooding, damage to crops, and increases in run-off/pollutants into coastal waters.</p> <p>A wetter climate may also lead to increases in vector-borne diseases (e.g., dengue).</p> <p>Impacts will be felt from periods of drought affecting human health, water supply and agriculture.</p>  |
| <b>Ocean acidification</b>           | <p>As atmospheric CO<sub>2</sub> concentrations continue to rise, oceans will warm and continue to acidify. Over the last two centuries in Palau, the aragonite saturation state (a proxy for coral reef growth rate) has declined from 4.5 to 3.9. Models suggest the aragonite saturation state will continue to decrease to 3.5 (marginal conditions to support coral reefs) by 2030 and decrease further to values where coral reefs have not historically been found (&lt; 3.0).</p> | <p>Ocean acidification (OA) affects many marine organisms that rely on calcium carbonate to build their shells/skeleton (e.g., corals, clams, mussels). OA can result in decreased growth and reproduction and weaker and more brittle skeletons, prone to increased damage from storms. Corals are critical because they provide habitats for fish, support food/livelihoods, income from tourism, medicines, and coastal protection to islands.</p> |
| <b>El Niño/La Niña – ENSO</b>        | <p>El Niño and La Niña events will continue to occur in the future, but there is little consensus on whether they will change in intensity or frequency.</p>  | <p>Impacts vary depending on ENSO years but affect sea level, precipitation patterns, temperature (air and sea), and storm patterns resulting in intense flooding and drought. In Palau, drought may occur in the year after a moderate/strong El Niño. ENSO also can impact coral reefs (e.g., through bleaching).</p>   |

**Acknowledgements:** Climate projections are derived using the Global Climate Model data from the Coupled Model Intercomparison Project, Phase 5 (CMIP5). Projections provided are for the greenhouse gas and aerosol concentration emission scenario RCP8.5 (Representative Concentration Pathway; very high emissions). Climate projection data are obtained from the Australian Bureau of Meteorology and CSIRO (2014). Climate Variability, Extremes and Change in the Western Tropical Pacific: New Science and Updated Country Reports and The Pacific Islands Regional Climate Assessment (PIRCA): <http://www.pacificrisa.org/projects/pirca/>